

## NY Breeding Bird Atlas III Breeding Codes

Every time you're out birding, you're observing bird behavior. It could be a song, some territorial squabble, or even a bird building a nest. Matching bird behaviors to specific codes is all you have to do to atlas!

Breeding behaviors are divided into four categories of breeding evidence: Observed, Possible, Probable, and Confirmed. The behavior codes below are listed from low to high. **Every time you are out atlasing, enter the highest breeding code observed for each species and be sure to submit your checklist in the atlas portal.** If you are in doubt about which code to use, please describe the behavior(s) you observed in the comments field for that species in your eBird checklist. Additional materials are provided on the atlas website ([ebird.org/atlasny](http://ebird.org/atlasny)) to help you determine when a species is likely to be breeding and appropriate codes for each species.

### — OBSERVED —

**No code, checklist in atlas portal:** Species present but without evidence of breeding and outside of suitable nesting habitat. No code is required. Any birds you enter on your checklists in the atlas portal without a breeding code are automatically 'Observed.'

**F – Flyover:** Birds flying high overhead or in direct flight. This is a behavior code that doesn't indicate breeding, therefore it is in the 'Observed' category. Tree Swallows foraging for insects are not considered a flyover because they are not in direct flight. Please see [the eBird help pages](#) for more detail.

### — POSSIBLE —

**H – In Appropriate Habitat:** Adult in suitable nesting habitat during its breeding season. Note that this code has both a habitat and breeding season component, which can be very different for residents vs migrants.

**S – Singing Bird:** Singing bird present in suitable nesting habitat. This code is also used for non-songbirds giving their primary vocalization, such as woodpecker drumming, owl calls, rail vocalizations, turkeys gobbling, and woodcock peenting. If in doubt, refer to the Acceptable Codes Chart. This code can be used for species where both males and females sing (e.g., Northern Cardinal, Bicknell's Thrush). If a bird is heard singing at the same place on a second trip during the breeding season, it may qualify as S7.

### — PROBABLE —

**S7 – Singing Bird Present 7+ Days:** Singing bird present at same location on at least two occasions 7 or more days apart. Both observations must be made by the same observer. Do not use if you observed the species singing a week earlier elsewhere in your block; this behavior presumes a permanent territory. Observations must be made within the same breeding season; observations from different years do not apply.

**M – Multiple (7+) Singing Birds:** Multiple (7 or more) singing or territorial birds of a species detected within a block within a single breeding season. If you are submitting separate checklists for different locations and habitats in your block (recommended) and you don't have all 7 birds on a single checklist, use this code on the checklist where you reach a total of 7 birds and make a note in the comment field that you are including birds from other checklists in the block.

**P – Pair in Suitable Habitat:** Pair observed in suitable nesting habitat. Only use this code when you are fairly certain that a mated pair of birds has been observed. Look for behavioral cues like sitting in close proximity to determine if you have a male-female pair, particularly in species where males and females look the same.

**T – Territorial Defense:** Permanent territory presumed through defense of breeding territory by fighting or chasing. While this is generally used for individuals of the *same species*, an interaction between members of different species may fall under this code when it appears to be territorial defense. Also see "A – Agitated Behavior." Because territoriality involves the defense of a fixed area, it may be useful to map locations of individuals to determine if they are defending the same general area when surveying the block on future visits.

**C – Courtship, Display, or Copulation:** Courtship behavior or copulation between a male and female. Courtship behavior includes transfer of food, displays, and grooming between a pair of birds.

**N – Visiting Probable Nest Site:** Repeated visits to a probable nest site. This is especially useful for cavity nesters or for a shrub-nesting species that flies into the same thicket and disappears on several occasions.

**A – Agitated Behavior:** Agitated behavior or anxiety calls from adults indicating a nest site or young in the vicinity. This code refers to a stronger reaction to intruders than those exhibited by “T – Territorial Defense,” usually against *other species*, such as brood parasites, nest predators, and humans. Do not use this code for agitation induced by “pishing” or playing recordings. This code also excludes mobbing behavior that species engage in year-round (e.g., mobbing an owl).

**B – Woodpecker/Wren Nest Building:** Nest-building (including carrying nesting material) by wrens or excavation of cavities by woodpeckers. Wrens may build “dummy” nests before the female selects a nest. Woodpeckers often drill holes for roosting.

— CONFIRMED —

**PE – Physiological Evidence:** Physiological evidence of breeding based on a bird in the hand. This code is used primarily by bird banders and includes evidence such as a highly vascularized swollen incubation (brood) patch, cloacal protuberance, or an egg in the oviduct. If possible, record the MAPS score for BP and CP.

**CN – Carrying Nesting Material:** Adult carrying nesting material to an unseen nest, such as sticks, grass, mud, and cobwebs. For raptors, be sure the material is not simply incidental to prey capture/transport. For wrens, use B.

**NB – Nest Building:** Nest-building observed at the actual nest site, excluding wrens and woodpeckers.

**DD – Distraction Display:** Distraction displays and injury feigning in attempt to draw intruder away from nest or young.

**UN – Used Nest:** Used nest found, but no adult birds seen nearby. Use *only if the nest was used during the Atlas period* (2020-2024) and you are *certain* of the species. Add comments detailing how you identified the nest. Do not collect the nest, but do take a photograph if possible. Enter ‘0’ if no individuals of that species are observed during your visit.

**ON – Occupied Nest:** Occupied nest indicated by adult sitting in nest in incubating position, adult entering nest site and remaining, or exchange of incubation duties by the pair. This code is useful for nests high in trees, on cliffs, and in chimneys where the contents of the nest and incubating or brooding adult cannot be seen.

**FL – Recently Fledged & Precocial Young:** Recently fledged or downy young still dependent upon adults and presumed incapable of extended flights from nest site. Look for retained downy feathers, a yellow gape, a short tail (shorter than the wings), clumsy flight and landings, and a bird incapable of feeding itself. Beware of family groups late in the breeding season which may still be interacting but are far from their breeding location. Young cowbirds begging for food confirm both the cowbird and the host species. If you find a dead fledgling and don’t see an adult of the same species, use a count of ‘0’ and enter the FL code.

**CF – Carrying Food:** Adult carrying food for young or incubating partner. Provide comments if used for corvids, raptors, terns, and other species that regularly carry food for courtship, caching, or other purposes. One of the best signs to look for is the repeated carrying of food in the same direction.

**FY – Feeding Young:** Adult bird feeding young that have left the nest. This code should only be used for recently fledged (juvenile) birds, not for immature birds that have left their breeding grounds (such as raptors and terns). Use the NY code for nestlings being fed by an adult.

**FS – Carrying Fecal Sac:** Adult carrying fecal sac or egg shell fragments. Many passerine adults keep their nests clean by carrying membranous, white fecal sacs and broken eggshells away from the nest. Note that only songbirds and woodpeckers produce fecal sacs and this code should only be applied to these groups of species. Eggshell fragments found on the ground should not be coded.

**NE – Nest with Eggs:** Nest with eggs. Be careful not to disturb the vicinity of the nest. Confirm the species by waiting at a distance until adult returns. If no birds are seen, and you are sure of the species, enter “0” for the count. If a cowbird egg is found in the nest, use code NE for both the cowbird and the host species; if no individual cowbirds were seen that day, then enter a “0” in the cowbird number field during data entry.

**NY – Nest with Young:** Nest with young seen or heard. Keep your distance so nestlings are not prematurely flushed from the nest. Include the nestlings in your species count. Presence of cowbird young confirms both the cowbird and the host species.