



Priority species are bolded (location, habitat, nest info). Sensitive species are asterisked (report to coordinator). Colonial species / Bald Eagle are italicized (nest location info only).



Nesting -----Migration / Dispersal




Priority species are bolded (location, habitat, nest info). Sensitive species are asterisked (report to coordinator). Colonial species / Bald Eagle are italicized (nest location info only).



Nesting -----Migration / Dispersal




Priority species are bolded (location, habitat, nest info). Sensitive species are asterisked (report to coordinator). Colonial species / Bald Eagle are italicized (nest location info only).





Priority species are bolded (location, habitat, nest info). Sensitive species are asterisked (report to coordinator). Colonial species / Bald Eagle are italicized (nest location info only).



Nesting -----Migration / Dispersal



Migration and dispersal dates were created using safe dates prepared for the Maryland & DC Breeding Bird Atlas 3.

Nesting dates were created using each species' first egg date + the last egg date + the mean incubation period + the mean nestling period.

First and last egg dates came from the *Yellowbook* (Iliff, Ringler, and Stasz) and/or the *2nd Atlas of the Breeding Birds of Maryland and the District of Columbia* (Ed. Walter G. Ellison). Where first or last egg dates were not provided by these sources, they were estimated from chick records in the *2nd Atlas of the Breeding Birds of Maryland and the District of Columbia*, provided by experts, or obtained from information found in Birds of the World (https://birdsoftheworld.org). If the mean incubation or nestling period was not available, the median was used; when two adjacent numbers were provided, the higher one was used.